

Assess the Severity of Restless Legs Syndrome

USING THE INTERNATIONAL RLS (IRLS) RATING SCALE

The IRLS Rating Scale was developed by the **International Restless Legs Syndrome Study Group (IRLSSG)** to assess the severity of a patient's RLS symptoms.¹ Ask your patient the 10 questions listed below and have them rate their symptoms from 0 to 4. Mark each answer and total their IRLS score. It will range from 0 to 40.

The IRLS Rating Scale was validated in a controlled study and found to have high levels of internal consistency, interexaminer reliability, and test-retest reliability.¹ The IRLSSG is an organization of professionals committed to advancing basic and clinical research on RLS. To learn more about the group, visit www.irlssg.org.

In the past week:

Score

Overall, how would you rate the RLS discomfort in your legs or arms?

4 Very severe 3 Severe 2 Moderate 1 Mild 0 None

Overall, how would you rate the need to move around because of your RLS symptoms?

4 Very severe 3 Severe 2 Moderate 1 Mild 0 None

Overall, how much relief of your RLS arm or leg discomfort did you get from moving around?

4 No relief 3 Mild relief 2 Moderate relief 1 Either complete or almost complete relief 0 No RLS symptoms to be relieved

How severe was your sleep disturbance from your RLS symptoms?

4 Very severe 3 Severe 2 Moderate 1 Mild 0 None

How severe was your tiredness or sleepiness during the day due to your RLS symptoms?

4 Very severe 3 Severe 2 Moderate 1 Mild 0 None

How severe was your RLS as a whole?

4 Very severe 3 Severe 2 Moderate 1 Mild 0 None

How often did you get RLS symptoms?

4 Very often (6–7 days in 1 week). 3 Often (4–5 days in 1 week). 2 Sometimes (2–3 days in 1 week). 1 Occasionally (1 day in 1 week). 0 Never

When you had RLS symptoms, how severe were they on average?

4 Very severe (8 h or more per 24 h) 3 Severe (3–8 h per 24 h) 2 Moderate (1–3 h per 24 h) 1 Mild (less than 1 h per 24 h) 0 None

Overall, how severe was the impact of your RLS symptoms, on your ability to carry out your daily affairs, for example, carrying out a satisfactory family, home, social, school, or work life?

4 Very severe 3 Severe 2 Moderate 1 Mild 0 None

How severe was your mood disturbance from your RLS symptoms—for example angry, depressed, sad, anxious, or irritable?

4 Very severe 3 Severe 2 Moderate 1 Mild 0 None

Add the scores from above and share the total with your patient.

Mild 0 to 10

Moderate 11 to 20

Severe 21 to 30

Very Severe 31 to 40

Total Score



Distributed by Arbor Pharmaceuticals with permission from the International Restless Legs Syndrome Study Group. This scale is copyrighted by the International Restless Legs Syndrome Study Group. ©IRLS Study Group 2013. All rights reserved.

For information about approved uses of this scale and about RLS, contact www.IRLSSG.org. IRLS contact information and permission to use: MAPI Research Trust, Lyon, France. Email: PROinformation@mapi-trust.org

Reference: 1. The International Restless Legs Syndrome Study Group. Validation of the International Restless Legs Syndrome Study Group Rating Scale for Restless Legs Syndrome. *Sleep Med.* 2003;4(2):121-132.

 **arbor**
PHARMACEUTICALS, LLC